

Uniqueness of Hamiltonian volume minimizing Lagrangian submanifolds which are Hamiltonian isotopic to $\mathbb{R}P^n$ in $\mathbb{C}P^n$ *

Hiroshi Iriyeh

1 Introduction

The equator on S^2 has the least length among all its images under area bisecting deformations. A symplectic higher dimensional generalization of this result was obtained by Y.-G. Oh [5] and B. Kleiner in 1990.

Theorem 1 (Kleiner-Oh). *The standard real form $\mathbb{R}P^n \subset \mathbb{C}P^n$ has the least volume among all its images under Hamiltonian isotopies.*

In other words, for any Hamiltonian diffeomorphism $\phi \in \text{Ham}(\mathbb{C}P^n)$ of the n -dimensional complex projective space $\mathbb{C}P^n$, we have

$$\text{vol}(\phi(\mathbb{R}P^n)) \geq \text{vol}(\mathbb{R}P^n).$$

A minimal (or Hamiltonian-minimal) Lagrangian submanifold with such a property is said to be *Hamiltonian volume minimizing*.

The purpose of this short note is to prove the uniqueness of Hamiltonian volume minimizing Lagrangian submanifolds which are Hamiltonian isotopic to $\mathbb{R}P^n$ in $\mathbb{C}P^n$ modulo isometric group actions. More precisely,

Theorem 2. *Let ϕ be a Hamiltonian diffeomorphism of $\mathbb{C}P^n$ with the standard Fubini-Study Kähler form ω . If $\text{vol}(\phi(\mathbb{R}P^n)) = \text{vol}(\mathbb{R}P^n)$ holds, then ϕ must be an isometry of $\mathbb{C}P^n$.*

Remark 3. *In the case of a totally geodesic Lagrangian torus in $S^2(1) \times S^2(1)$, an analogous result of the above theorems was proved in [3] and [4].*

The author is grateful to Professor T. Otofui for helpful comments.

*2000 Mathematics Subject Classification. Primary 53C40; Secondary 53C65.

2 Review on Kleiner-Oh's method

In this section, we review a proof of theorem 1 by Kleiner and Oh. This is a combination of the Lagrangian intersection theorem in symplectic geometry and Poincaré formula in integral geometry. Let $(\mathbb{C}P^n, \omega, J)$ be the n -dimensional complex projective space with the Fubini-Study Kähler form ω and the standard complex structure J .

Theorem 4 (Givental [1]). *For any Hamiltonian diffeomorphism ϕ of $\mathbb{C}P^n$ such that $\mathbb{R}P^n$ and $\phi(\mathbb{R}P^n)$ intersect transversely, the inequality*

$$\sharp(\mathbb{R}P^n \cap \phi(\mathbb{R}P^n)) \geq n + 1 \quad (1)$$

holds.

Next, we explain the Poincaré formula for Lagrangian submanifolds in $\mathbb{C}P^n$.

Theorem 5 (Kleiner/Howard [2]). *Let $\mathbb{C}P^n = U(n+1)/U(1) \times U(n)$ be the complex projective space. Let L_1 and L_2 be compact Lagrangian submanifolds in $\mathbb{C}P^n$. Then we have*

$$\int_{U(n+1)} \sharp(L_1 \cap gL_2) d\mu(g) = C \text{vol}(L_1) \text{vol}(L_2) \quad (2)$$

where

$$C = \frac{(n+1) \text{vol}(U(n+1))}{\text{vol}(\mathbb{R}P^n)^2}.$$

These two theorems yield the Hamiltonian volume minimizing property of $\mathbb{R}P^n$ as follows. By equation (2), we have

$$\begin{aligned} & \frac{(n+1) \text{vol}(U(n+1))}{\text{vol}(\mathbb{R}P^n)^2} \text{vol}(\mathbb{R}P^n) \text{vol}(\phi(\mathbb{R}P^n)) \\ &= \int_{U(n+1)} \sharp(\mathbb{R}P^n \cap g\phi(\mathbb{R}P^n)) d\mu(g). \end{aligned} \quad (3)$$

Since $g \in U(n+1)$ is a Hamiltonian diffeomorphism, by the group structure of $\text{Ham}(\mathbb{C}P^n)$, $g\phi$ is also a Hamiltonian diffeomorphism. By equation (1), we have

$$\text{RHS of (3)} \geq \int_{U(n+1)} (n+1) d\mu(g) = (n+1) \text{vol}(U(n+1)). \quad (4)$$

Hence,

$$\text{vol}(\phi(\mathbb{R}P^n)) \geq \text{vol}(\mathbb{R}P^n).$$

3 Proof

First of all, we exhibit the key fact to prove theorem 2.

Theorem 6 (Oh [6]). *Let L be an embedded closed Lagrangian submanifold in $\mathbb{C}P^n$. If L satisfies that*

$$\sharp(L \cap gL) = SB(L, \mathbb{Z}_2)$$

for all $g \in U(n+1)$ such that L and gL intersect transversely, then L is the standard totally geodesic $\mathbb{R}P^n$ in $\mathbb{C}P^n$. Here, $SB(L, \mathbb{Z}_2)$ denotes the sum of $\mathbb{Z}_2$ -Betti numbers of L .

Assume that $\text{vol}(\phi(\mathbb{R}P^n)) = \text{vol}(\mathbb{R}P^n)$ for $\phi \in \text{Ham}(\mathbb{C}P^n)$. Then we have

$$\int_{U(n+1)} \sharp(\mathbb{R}P^n \cap g\phi(\mathbb{R}P^n)) d\mu(g) = \int_{U(n+1)} (n+1) d\mu(g). \quad (5)$$

On the other hand, by assumption,

$$\begin{aligned} & \int_{U(n+1)} \sharp(\phi(\mathbb{R}P^n) \cap g\phi(\mathbb{R}P^n)) d\mu(g) \\ &= C \text{vol}(\phi(\mathbb{R}P^n)) \text{vol}(\phi(\mathbb{R}P^n)) \\ &= C \text{vol}(\mathbb{R}P^n) \text{vol}(\phi(\mathbb{R}P^n)) \\ &= \int_{U(n+1)} \sharp(\mathbb{R}P^n \cap g\phi(\mathbb{R}P^n)) d\mu(g). \end{aligned} \quad (6)$$

By equations (5) and (6), we have

$$\int_{U(n+1)} \{\sharp(\phi(\mathbb{R}P^n) \cap g\phi(\mathbb{R}P^n)) - (n+1)\} d\mu(g) = 0.$$

By theorem 4, for almost every $g \in U(n+1)$, we have

$$\sharp(\phi(\mathbb{R}P^n) \cap g\phi(\mathbb{R}P^n)) = n+1. \quad (7)$$

Take $g \in U(n+1)$ which satisfies that $\phi(\mathbb{R}P^n)$ and $g\phi(\mathbb{R}P^n)$ intersect transversely. If $\sharp(\phi(\mathbb{R}P^n) \cap g\phi(\mathbb{R}P^n)) > n+1$, then this inequality holds for all $g' \in U(n+1)$ near g . This contradicts equation (7). Hence, $\sharp(\phi(\mathbb{R}P^n) \cap g\phi(\mathbb{R}P^n)) = n+1$ for all $g \in U(n+1)$ such that $\phi(\mathbb{R}P^n)$ and $g\phi(\mathbb{R}P^n)$ intersect transversely.

By theorem 6, the Lagrangian submanifold $\phi(\mathbb{R}P^n)$ must be the standard totally geodesic $\mathbb{R}P^n \subset \mathbb{C}P^n$. Therefore, $\phi \in U(n+1)$. $\square$

References

- [1] A. B. Givental, *Periodic maps in symplectic topology*, Funct. Anal. Appl., **23** (1989), 37-52.
- [2] R. Howard, *The kinematic formula in Riemannian homogeneous spaces*, Mem. Amer. Math. Soc., No.509, **106** (1993).
- [3] H. Iriyeh, H. Ono and T. Sakai, *Integral geometry and Hamiltonian volume minimizing property of a totally geodesic Lagrangian torus in $S^2 \times S^2$* , Proc. Japan Acad., **79**, Ser. A (2003), 167–170.
- [4] H. Iriyeh, H. Ono and T. Sakai, *Hamiltonian volume minimizing property of Lagrangian submanifolds*, RIMS Kokyuroku, **1403** (2004), 10–17.
- [5] Y.-G. Oh, *Second variation and stabilities of minimal lagrangian submanifolds in Kähler manifolds*, Invent. Math. **101** (1990), 501-519.
- [6] Y.-G. Oh, *Tight Lagrangian submanifolds in $\mathbb{C}P^n$* , Math. Z. **207** (1991), 409-416.

SCHOOL OF ENGINEERING
TOKYO DENKI UNIVERSITY
KANDA-NISHIKI-CHO, CHIYODA-KU
TOKYO, 101-8457
JAPAN
e-mail : hirie@im.dendai.ac.jp